arXiv:1703.08362v1 [cs.IT] 24 Mar 2017

A new class of three-weight linear codes from weakly regular plateaued functions ⋆

Sihem Mesnager[1,2,3], Ferruh Özbudak[4,5], and Ahmet Sınak[2,5,6]

[1] Department of Mathematics, University of Paris VIII, France
[2] LAGA, UMR 7539, CNRS, University of Paris VIII and University of Paris XIII, France
[3] Telecom ParisTech, France
[4] Department of Mathematics, Middle East Technical University, Turkey
[5] Institute of Applied Mathematics, Middle East Technical University, Turkey
[6] Department of Mathematics and Computer Sciences, Necmettin Erbakan University, Turkey
smesnager@univ-paris8.fr,{ozbudak,sahmet}@metu.edu.tr

Abstract. Linear codes with few weights have many applications in secret sharing schemes, authentication codes, communication and strongly regular graphs. In this paper, we consider linear codes with three weights in arbitrary characteristic. To do this, we generalize the recent contribution of Mesnager given in [Cryptography and Communications 9(1), 71-84, 2017]. We first present a new class of binary linear codes with three weights from plateaued Boolean functions and their weight distributions. We next introduce the notion of (weakly) regular plateaued functions in odd characteristic p and give concrete examples of these functions. Moreover, we construct a new class of three-weight linear p-ary codes from weakly regular plateaued functions and determine their weight distributions. We finally analyse the constructed linear codes for secret sharing schemes.

Keywords: Binary codes, linear codes, p-ary codes, p-ary functions, secret sharing schemes, weakly regular plateaued, weight distribution.

1 Introduction

Error correcting codes have many applications in communication systems, data storage devices and consumer electronics. The construction of linear codes with few weights has been widely studied (see, e.g., [3,4,10,12,14,16]) since these codes have many applications in consumer electronics, secret sharing schemes, authentication codes, communication, data storage system, association schemes, and strongly regular graphs. Recently, in [2], Ding has published a valuable survey on the construction of binary linear codes from Boolean functions. The notion of plateaued Boolean functions, as an extension of the notion of bent Boolean functions, has been introduced in [15] by Zheng and Zhang (1999), and then generalized to arbitrary characteristic: the so-called p-ary plateaued functions from $\mathbb{F}_{p^n}$ to $\mathbb{F}_p$ (see, e.g., [11]). Several researchers have studied plateaued functions since they have many applications in cryptography, sequence theory and coding theory. In particular, p-ary bent functions (mostly, quadratic and weakly regular bent functions) have been used in coding theory to construct linear codes with few weights. Very recently, Mesnager [10] has constructed a new family of three-weight linear codes from weakly regular bent functions in arbitrary characteristic based on a generic construction. Within this framework, the aim of this paper is to construct a class of linear codes with few weights from weakly regular plateaued functions in arbitrary characteristic and determine their weight distributions.

⋆ The Extended Abstract of this work was submitted to WCC-2017 (the Tenth International Workshop on Coding and Cryptography).

The paper is structured as follows. Section 2 sets the main notations and recalls some basic results in coding theory and number theory. In Section 3, we introduce the notion of (weakly) regular plateaued functions in odd characteristic p. We then give concrete examples to show the existence of (weakly) regular plateaued p-ary functions. Section 4 constructs a new class of three-weight linear p-ary (resp. binary) codes from weakly regular p-ary plateaued (resp. plateaued Boolean) functions based on a generic construction. We also determine the weight distributions of the constructed linear codes in this paper. Finally, in Section 5, we observe that all nonzero codewords of the constructed linear codes are minimal for almost all cases.

2 Preliminaries

In this section, we set main notations and give some basic results on p-ary functions, coding theory and number theory, which will be used in the sequel.
For any set E, $\#E$ denotes the cardinality of E and $E^\star = E \setminus \{0\}$. Given a complex number $z \in \mathbb{C}$, $|z|$ denotes the absolute value of z, where $\mathbb{C}$ is the field of complex numbers. Let $\mathbb{F}_{p^m}$ be the finite field with p^m elements, where p is a prime and $m \geq 1$ is a positive integer. Then, $\mathbb{F}_{p^m}^\star = \langle \zeta \rangle$ is a multiplicative cyclic group of order $p^m - 1$ with generator ζ, and $\mathbb{F}_p$ is the prime field of $\mathbb{F}_{p^m}$. The extension field $\mathbb{F}_{p^m}$ can be seen as an m-dimensional vector space over $\mathbb{F}_p$, denoted by $\mathbb{F}_p^m$. The absolute trace function $\mathrm{Tr}_p^{p^m} : \mathbb{F}_{p^m} \to \mathbb{F}_p$ is defined as $\mathrm{Tr}_p^{p^m}(x) := \sum_{i=0}^{m-1} x^{p^i}$. Recall that $\mathrm{Tr}_p^{p^m}$ is $\mathbb{F}_p$-linear. Given a function $f : \mathbb{F}_{p^m} \longrightarrow \mathbb{F}_p$, the direct and inverse Walsh transform of f are defined, respectively, by:

$$\widehat{\chi_f}(b) = \sum_{x \in \mathbb{F}_{p^m}} \xi_p^{f(x) - \mathrm{Tr}_p^{p^m}(bx)} \text{ and}$$

$$\xi_p^{f(x)} = \frac{1}{p^m} \sum_{b \in \mathbb{F}_{p^m}} \widehat{\chi_f}(b) \xi_p^{\mathrm{Tr}_p^{p^m}(bx)}, \tag{1}$$

where $\xi_p = e^{\frac{2\pi\sqrt{-1}}{p}}$ is a primitive p-th root of unity. The set $\{b \in \mathbb{F}_{p^m} : \widehat{\chi_f}(b) \neq 0\}$ is called the Walsh support of f, and is denoted by $Supp\,(\widehat{\chi_f})$. For a nonnegative integer i, the moment of Walsh transform of f is defined by $S_i(f) = \sum_{b \in \mathbb{F}_{p^m}} |\widehat{\chi_f}(b)|^{2i}$ with the convention $S_0(f) = p^m$, and $S_1(f) = p^{2m}$ is known as the *Parseval identity*. Recall that f is said to be *balanced* over $\mathbb{F}_p$ if $\#\{x \in \mathbb{F}_p^n : f(x) = k\} = p^{m-1}$ for each $k \in \mathbb{F}_p$, i.e., f takes every value of $\mathbb{F}_p$ the same number p^{m-1} times; otherwise, it is called *unbalanced*.

Basic background in number theory. We now recall the basic facts of the Legendre symbol and cyclotomic field. Let a be a positive integer and p be an odd prime. We say that a is a quadratic residue modulo p if $\sqrt{a} \in \mathbb{F}_p^\star$, and a is a quadratic non-residue modulo p if $\sqrt{a} \notin \mathbb{F}_p^\star$. The *Legendre symbol* is defined as

$$\left(\frac{a}{p}\right) := \begin{cases} 0 & \text{if } p \mid a, \\ 1 & \text{if } a \text{ is a quadratic residue modulo } p, \\ -1 & \text{if } a \text{ is a quadratic non-residue modulo } p. \end{cases}$$

The Legendre symbol satisfies $\left(\frac{a}{p}\right) \equiv a^{\frac{p-1}{2}} \pmod p$, and

$$\left(\frac{-1}{p}\right) \equiv (-1)^{\frac{p-1}{2}} \pmod p = \begin{cases} 1 & \text{if } p \equiv 1 \pmod 4, \\ -1 & \text{if } p \equiv 3 \pmod 4. \end{cases} \tag{2}$$

Throughout this paper, p^* denotes $\left(\frac{-1}{p}\right) p$, $\left(\frac{a}{p}\right)$ denotes the Legendre symbol for $a \in \mathbb{F}_p^\star$, $\mathbb{Z}$ is the rational integer ring and $\mathbb{Q}$ is the rational field. The ring of integers in $\mathbb{Q}(\xi_p)$ is $\mathcal{O}_K := \mathbb{Z}(\xi_p)$. An integral basis of $\mathcal{O}_{\mathbb{Q}(\xi_p)}$ is the set $\{\xi_p^i \mid 1 \le i \le p-1\}$. The Galois field extension $\mathbb{Q}(\xi_p)/\mathbb{Q}$ of degree $p-1$ is the Galois group $Gal(\mathbb{Q}(\xi_p)/\mathbb{Q}) = \{\sigma_a \mid a \in (\mathbb{Z}/p\mathbb{Z})^\star\}$, where the automorphism σ_a of $\mathbb{Q}(\xi_p)$ is defined by $\sigma_a(\xi_p) = \xi_p^a$. The field $\mathbb{Q}(\xi_p)$ has a unique quadratic subfield $\mathbb{Q}(\sqrt{p^*})$. For $a \in \mathbb{F}_p^\star$, we have $\sigma_a(\sqrt{p^*}) = \left(\frac{a}{p}\right)\sqrt{p^*}$. Hence, the Galois group $Gal(\mathbb{Q}(\sqrt{p^*})/\mathbb{Q} = \{1, \sigma_\gamma\}$ for any $\gamma \in \mathbb{F}_p$ such that $\sqrt{\gamma} \notin \mathbb{F}_p^\star$. The reader is referred to [13] for further reading on cyclotomic fields.

Basic background in coding theory. Let q be a prime power and n be a positive integer. The support of a vector $\tilde{a} = (a_0, \ldots, a_{n-1}) \in \mathbb{F}_q^n$ is defined as $supp(\tilde{a}) := \{0 \le i \le n-1 : a_i \neq 0\}$. The Hamming weight of $\tilde{a} \in \mathbb{F}_q^n$, denoted by $wt(\tilde{a})$, is the cardinality of its support, i.e., $wt(\tilde{a}) := \#supp(\tilde{a})$. A linear $[n,k]_q$ code $\mathcal{C}$ over $\mathbb{F}_q$ is a k-dimensional subspace of $\mathbb{F}_q^n$. A linear $[n,k,d]_q$ code $\mathcal{C}$ over $\mathbb{F}_q$ is a k-dimensional subspace of $\mathbb{F}_q^n$ with minimum Hamming distance d. The dual code of $\mathcal{C}$ is the linear code with parameters $[n, n-k, d^\perp]_q$ defined by

$$\mathcal{C}^\perp = \{\tilde{b} \in \mathbb{F}_q^n : \tilde{b} \cdot \tilde{a} = \tilde{0} \text{ for all } \tilde{a} \in \mathcal{C}\},$$

where " $\cdot$ " is an inner product in $\mathbb{F}_q^n$. Let A_w denote the number of codewords with Hamming weight w in $\mathcal{C}$ of length n. Then, $(1, A_1, \ldots, A_n)$ is the weight distribution of $\mathcal{C}$ and the polynomial $1 + A_1 y + \cdots + A_n y^n$ is called the weight enumerator of $\mathcal{C}$. The code $\mathcal{C}$ is called a t-weight code if the number of nonzero A_w in the weight distribution is t. For further reading on coding theory, we send the reader to [7].

3 On (weakly) regular plateaued p-ary functions

In this section, we introduce the notion of (weakly) regular plateaued functions in odd characteristic p and give some properties of these functions. We first recall the notion of plateaued functions.

Let $f : \mathbb{F}_{p^m} \longrightarrow \mathbb{F}_p$ be a function. A p-ary function f is called *bent* if all of its Walsh transform coefficients satisfy $|\widehat{\chi_f}(b)|^2 = p^m$, and *$r$-plateaued* if all of its Walsh transform coefficients satisfy $|\widehat{\chi_f}(b)|^2 \in \{0, p^{m+r}\}$, where r is an integer with $0 \le r \le m$. We point out that a 0-plateaued function is bent. In characteristic 2, it is safe to say that f is *r-plateaued Boolean function* if $\widehat{\chi_f}(b) \in \{0, \pm 2^{(m+r)/2}\}$ for all $b \in \mathbb{F}_{2^m}$. By the Parseval identity, we have (see, e.g., [11]):

Lemma 1. *Let p be any prime and $f : \mathbb{F}_{p^m} \to \mathbb{F}_p$ be r-plateaued. Then for $b \in \mathbb{F}_{p^m}$, $|\widehat{\chi_f}(b)|^2$ takes p^{m-r} times the value p^{m+r} and $p^m - p^{m-r}$ times the value 0.*

Lemma 2. *Let $f : \mathbb{F}_{2^m} \to \mathbb{F}_2$ be a r-plateaued Boolean function. Then for $b \in \mathbb{F}_{2^m}$, the Walsh distribution of f is given by*

$$\widehat{\chi_f}(b) = \begin{cases} 2^{\frac{m+r}{2}}, & 2^{m-r-1} + 2^{\frac{m-r-2}{2}} \text{ times,} \\ 0, & 2^m - 2^{m-r} \text{ times,} \\ -2^{\frac{m+r}{2}}, & 2^{m-r-1} - 2^{\frac{m-r-2}{2}} \text{ times.} \end{cases}$$

We recall the notion of (weakly) regular bent functions in odd characteristic p (see, e.g., [6]). For an odd prime p, the Walsh transform coefficients of a p-ary bent function f satisfy

$$\widehat{\chi_f}(b) = \begin{cases} \pm p^{\frac{m}{2}} \xi_p^{f^\star(b)}, & \text{if } m \text{ is even or } m \text{ is odd and } p \equiv 1 \pmod 4, \\ \pm i p^{\frac{m}{2}} \xi_p^{f^\star(b)}, & \text{if } m \text{ is odd and } p \equiv 3 \pmod 4, \end{cases}$$

where i is a complex primitive 4-th root of unity and f^* is called the dual of f. A bent function f is called *regular* if for all $b \in \mathbb{F}_{p^m}$, $\widehat{\chi_f}(b) = p^{\frac{m}{2}} \xi_p^{f^*(b)}$, and *weakly regular* if there exists a complex number u having unit magnitude (in fact, $|u| = 1$ and u does not depend on b) such that $\widehat{\chi_f}(b) = u p^{\frac{m}{2}} \xi_p^{f^*(b)}$ for all $b \in \mathbb{F}_{p^m}$, where f^* is the dual of f; otherwise, f is called *non-weakly regular*.
Very recently, Hyun et al. [8] have proved that the Walsh transform coefficients of a p-ary r-plateaued function f satisfy

$$\widehat{\chi_f}(b) = \begin{cases} \pm p^{\frac{m+r}{2}} \xi_p^{g(b)}, 0 & \text{if } m+r \text{ is even or } m+r \text{ is odd and } p \equiv 1 \pmod 4, \\ \pm i p^{\frac{m+r}{2}} \xi_p^{g(b)}, 0 & \text{if } m+r \text{ is odd and } p \equiv 3 \pmod 4, \end{cases} \tag{3}$$

where i is a complex primitive 4-th root of unity and g is a p-ary function over $\mathbb{F}_{p^m}$ with $g(b) = 0$ for $b \notin Supp(\widehat{\chi_f})$. Notice that by definition of $g : \mathbb{F}_{p^m} \to \mathbb{F}_p$, it can be regarded as a mapping from $Supp(\widehat{\chi_f})$ to $\mathbb{F}_p$ since we have $g(b) = 0$ for all $b \notin Supp(\widehat{\chi_f})$.
The notion of weak regularity is meaningful for plateaued functions. We now introduce the notion of (weakly) regular plateaued functions, which covers a non-trivial subclass of the class of plateaued functions.

Definition 1. *Let p be an odd prime and $f : \mathbb{F}_{p^m} \to \mathbb{F}_p$ be a p-ary r-plateaued function, where r is an integer with $0 \le r \le m$. Then, f is called* regular p-ary r-plateaued *if $\widehat{\chi_f}(b) \in \{0, p^{\frac{m+r}{2}} \xi_p^{g(b)}\}$ for all $b \in \mathbb{F}_{p^m}$, where g is a p-ary function over $\mathbb{F}_{p^m}$ with $g(b) = 0$ for all $b \notin Supp(\widehat{\chi_f})$. Moreover, f is called* weakly regular p-ary r-plateaued *if there exists a complex number u having unit magnitude (that is, $|u| = 1$ and u does not depend on b) such that*

$$\widehat{\chi_f}(b) \in \left\{0, u p^{\frac{m+r}{2}} \xi_p^{g(b)}\right\}$$

for all $b \in \mathbb{F}_{p^m}$, where g is a p-ary function over $\mathbb{F}_{p^m}$ with $g(b) = 0$ for all $b \notin Supp(\widehat{\chi_f})$; otherwise, f is called non-weakly regular p-ary r-plateaued.

Notice that we have $\widehat{\chi_f}(b) = 0$ if $b \notin Supp(\widehat{\chi_f})$. Then it is safe to say that f is regular r-plateaued if $\widehat{\chi_f}(b) = p^{\frac{m+r}{2}} \xi_p^{g(b)}$ for all $b \in Supp(\widehat{\chi_f})$, and f is weakly regular r-plateaued if there exists a complex number u having unit magnitude such that

$$\widehat{\chi_f}(b) = u p^{\frac{m+r}{2}} \xi_p^{g(b)} \tag{4}$$

for all $b \in Supp(\widehat{\chi_f})$, where $|u| = 1$ (in fact, u can only be equal to ± 1 or $\pm i$ and it does not depend on b) and g is a p-ary function over $Supp(\widehat{\chi_f})$. By (3), regular r-plateaued functions can only exist for even $m+r$ and for odd $m+r$ with $p \equiv 1 \pmod 4$. We can derive from (4) the following result.

Lemma 3. *Let f be a weakly regular r-plateaued p-ary function. For all $b \in Supp(\widehat{\chi_f})$, we can say $\widehat{\chi_f}(b) = \epsilon\sqrt{p^*}^{m+r}\xi_p^{g(b)}$, where $\epsilon = \pm 1$ is the sign of $\widehat{\chi_f}$, p^* denotes $\left(\frac{-1}{p}\right)p$ and g is a p-ary function over $Supp(\widehat{\chi_f})$.*

Proof. By (2) and (3), using the fact that u does not depend on b in (4), we obtain the following:

If $m+r$ is even or $m+r$ is odd and $p \equiv 1 \pmod 4$, then $\left(\frac{-1}{p}\right)^{m+r} = 1$ and $u = \pm 1$ in (4). Hence, we have $\epsilon\sqrt{p^*}^{m+r} = \epsilon\sqrt{1}\sqrt{p}^{m+r} = u\sqrt{p}^{m+r}$, where $\epsilon = \pm 1$.

If $m+r$ is odd and $p \equiv 3 \pmod 4$, then $\left(\frac{-1}{p}\right) = -1$ and $u = \epsilon i$ in (4), where $\epsilon = \pm 1$. Hence, $\epsilon\sqrt{p^*}^{m+r} = \epsilon\sqrt{-1}^{m+r}\sqrt{p}^{m+r} = \epsilon i^{m+r}\sqrt{p}^{m+r} = \epsilon i\sqrt{p}^{m+r} = u\sqrt{p}^{m+r}$. The result now follows. □

Remark 1. Notice that the notion of (weakly) regular 0-plateaued functions coincides with the one of (weakly) regular bent functions. Indeed, if we have $|\widehat{\chi_f}(b)|^2 \in \{0, p^m\}$ for all $b \in \mathbb{F}_{p^m}$, then by the Parseval identity, $p^{2m} = p^m \# Supp(\widehat{\chi_f})$, and so, $\# Supp(\widehat{\chi_f}) = p^m$. Hence, a (weakly) regular 0-plateaued function is the (weakly) regular bent.

By MAGMA, we obtain several (weakly) regular r-plateaued functions, two of which are given as follows for $p = n = 3$.

Example 1. A function $f(x) = \mathrm{Tr}_3^{3^3}(\zeta^5 x^{11} + \zeta^{20}x^5 + \zeta^{11}x^4 + \zeta^2 x^3 + \zeta x^2)$ where $\mathbb{F}_{3^3}^\star = \langle\zeta\rangle$ with $\zeta^3 + 2\zeta + 1 = 0$ is regular 3-ary 1-plateaued with $\widehat{\chi_f}(b) \in \{0, 9\xi_3^{g(b)}\}$, where g is an unbalanced 3-ary function.

Example 2. A function $f(x) = \mathrm{Tr}_3^{3^3}(\zeta x^{13} + \zeta^7 x^4 + \zeta^7 x^3 + \zeta x^2)$ where $\mathbb{F}_{3^3}^\star = \langle\zeta\rangle$ with $\zeta^3 + 2\zeta + 1 = 0$ is weakly regular 3-ary 1-plateaued with $\widehat{\chi_f}(b) \in \{0, -9\xi_3^{g(b)}\}$, where g is an unbalanced 3-ary function. On the other hand, a function $\mathrm{Tr}_3^{3^3}(\zeta^{16}x^{13} + \zeta^2 x^4 + \zeta^2 x^3 + \zeta x^2)$ is non-weakly regular 3-ary 2-plateaued.

The following lemma will be used to determine the weight distributions of the constructed linear codes.

Lemma 4. *Let f be a weakly regular r-plateaued p-ary function, that is, for all $b \in Supp(\widehat{\chi_f})$ we have $\widehat{\chi_f}(b) = up^{\frac{m+r}{2}}\xi_p^{g(b)}$, where $|u| = 1$. Then, we have*

$$\widehat{\chi_g}(x) = u^{-1}p^{\frac{m-r}{2}}\xi_p^{f(-x)}.$$

Proof. By the inverse Walsh transform in (1), we have

$$\begin{aligned} u^{-1}p^{\frac{m+r}{2}}\xi_p^{f(x)} &= u^{-1}p^{\frac{m+r}{2}}\frac{1}{p^m}\sum_{b\in\mathbb{F}_{p^m}}\widehat{\chi_f}(b)\xi_p^{\mathrm{Tr}_p^{p^m}(bx)} \\ &= u^{-1}p^{\frac{m+r}{2}}\frac{1}{p^m}\sum_{b\in Supp(\widehat{\chi_f})}up^{\frac{m+r}{2}}\xi_p^{g(b)}\xi_p^{\mathrm{Tr}_p^{p^m}(bx)} \\ &= p^r\sum_{b\in Supp(\widehat{\chi_f})}\xi_p^{g(b)+\mathrm{Tr}_p^{p^m}(bx)} = p^r\widehat{\chi_g}(-x). \end{aligned}$$

□

4 A new class of three-weight linear codes from weakly regular plateaued functions

In this section, we construct a new class of linear codes with few weights from plateaued functions in arbitrary characteristic and determine their weight distributions (we shall analyse separately the binary case and the case when p is odd). For any $\alpha, \beta \in \mathbb{F}_{p^m}$, one can define a function

$$\begin{aligned} f_{\alpha,\beta} : \mathbb{F}_{p^m} &\longrightarrow \mathbb{F}_p \\ x &\longmapsto f_{\alpha,\beta}(x) := \mathrm{Tr}_p^{p^m}(\alpha\Psi(x) - \beta x), \end{aligned}$$

where Ψ is a polynomial from $\mathbb{F}_{p^m}$ to $\mathbb{F}_{p^m}$ such that $\Psi(0) = 0$. Then one can define a linear code $\mathcal{C}_\Psi$ of length $p^m - 1$ over $\mathbb{F}_p$ as:

$$\mathcal{C}_\Psi := \{\tilde{c}_{\alpha,\beta} = (f_{\alpha,\beta}(\zeta_1), f_{\alpha,\beta}(\zeta_2), \ldots, f_{\alpha,\beta}(\zeta_{p^m-1})) \mid \alpha, \beta \in \mathbb{F}_{p^m}\},$$

where $\zeta_1, \ldots, \zeta_{p^m-1}$ are the elements of $\mathbb{F}_{p^m}^\star$. In this context, the following main results have been obtained in [10] by Mesnager.

Proposition 1. *Let ψ_a be a function from $\mathbb{F}_{p^m}$ to $\mathbb{F}_p$ defined by $\psi_a(x) = \mathrm{Tr}_p^{p^m}(a\Psi(x))$, where $a \in \mathbb{F}_{p^m}$ and $\Psi : \mathbb{F}_{p^m} \to \mathbb{F}_{p^m}$ with $\Psi(0) = 0$. For all $\alpha, \beta \in \mathbb{F}_{p^m}$, we have*

$$wt(\tilde{c}_{\alpha,\beta}) = p^m - \frac{1}{p} \sum_{\omega \in \mathbb{F}_p} \widehat{\chi_{\psi_{\omega\alpha}}}(\omega\beta).$$

We are going to consider a subclass of the class of linear codes $\mathcal{C}_\Psi$. We assume $a = 1$ and $\alpha \in \mathbb{F}_p$. Then, we have $f_{\alpha,\beta}(x) = \alpha\psi_1(x) - \mathrm{Tr}_p^{p^m}(\beta x)$ and define a subcode $\mathcal{C}$ of $\mathcal{C}_\Psi$ as follows:

$$\mathcal{C} := \{\tilde{c}_{\alpha,\beta} = (f_{\alpha,\beta}(\zeta_1), f_{\alpha,\beta}(\zeta_2), \ldots, f_{\alpha,\beta}(\zeta_{p^m-1})) \mid \alpha \in \mathbb{F}_p, \beta \in \mathbb{F}_{p^m}\}, \tag{5}$$

where $\zeta_1, \ldots, \zeta_{p^m-1}$ are the elements of $\mathbb{F}_{p^m}^\star$. Then, a linear code $\mathcal{C}$ over $\mathbb{F}_p$ defined by (5) is a k-dimensional subspace of $\mathbb{F}_p^n$, where $k = m + 1$ and $n = p^m - 1$, and it is denoted by $[p^m - 1, m + 1]_p$. By Proposition 1, the Hamming weights of the codewords of $\mathcal{C}$ can be given as follows.

Proposition 2. *We keep the above arguments. For $\tilde{c}_{\alpha,\beta} \in \mathcal{C}$, if $\alpha = 0$, we have $wt(\tilde{c}_{0,0}) = 0$ and $wt(\tilde{c}_{0,\beta}) = p^m - p^{m-1}$ for $\beta \neq 0$, if $\alpha \in \mathbb{F}_p^\star$, for all $\beta \in \mathbb{F}_{p^m}$ we have*

$$wt(\tilde{c}_{\alpha,\beta}) = p^m - p^{m-1} - \frac{1}{p} \sum_{\omega \in \mathbb{F}_p^\star} \sigma_\omega \left(\sigma_\alpha(\widehat{\chi_{\psi_1}}(\alpha^{-1}\beta)) \right),$$

where α^{-1} is the multiplicative inverse of α in $\mathbb{F}_p^\star$ and σ_a is the automorphism of $\mathbb{Q}(\xi_p)$ for $a \in \mathbb{F}_p^\star$.

4.1 A new class of binary three-weight linear codes from plateaued Boolean functions

In this subsection, we present a new class of binary linear codes with few weights and their weight distributions using plateaued Boolean functions.
Let $p = 2$ and assume that $\psi_1(x) = \mathrm{Tr}_2^{2^m}(\Psi(x))$ is a r-plateaued Boolean function, where $m + r$ is even. For $\alpha \in \mathbb{F}_2$ and $\beta \in \mathbb{F}_{2^m}$, we compute the Hamming weights of the codewords and weight distribution of $\mathcal{C}$ defined by (5). By Proposition 2, if $\alpha = 0$, we have $wt(\tilde{c}_{0,0}) = 0$ and $wt(\tilde{c}_{0,\beta}) = 2^{m-1}$ for $\beta \neq 0$, if $\alpha = 1$ and $\beta \in \mathbb{F}_{2^m}$, we have $wt(\tilde{c}_{1,\beta}) = 2^{m-1} - \frac{1}{2}\widehat{\chi_{\psi_1}}(\beta)$. By Lemma 2, we have for all $\beta \in \mathbb{F}_{2^m}$,

$$wt(\tilde{c}_{1,\beta}) = \begin{cases} 2^{m-1} - 2^{\frac{m+r-2}{2}}, 2^{m-r-1} + 2^{\frac{m-r-2}{2}} \text{ times}, \\ 2^{m-1}, \qquad\qquad 2^m - 2^{m-r} \text{ times}, \\ 2^{m-1} + 2^{\frac{m+r-2}{2}}, 2^{m-r-1} - 2^{\frac{m-r-2}{2}} \text{ times}. \end{cases}$$

We give in the following theorem the Hamming weights of the codewords and the weight distribution of $\mathcal{C}$.

Theorem 1. *Let $p = 2$ and $\mathcal{C}$ be a binary linear $[2^m - 1, m + 1]$ code defined by (5). Assume that ψ_1 is a r-plateaued Boolean function, where $m + r$ is even with $0 \leq r \leq m - 2$ for $2 \leq m$. Then, the Hamming weight of codewords and the weight distribution of $\mathcal{C}$ are as in Table 1.*

Hamming weight w	Multiplicity A_w
0	1
2^{m-1}	$2^{m+1} - 2^{m-r} - 1$
$2^{m-1} - 2^{\frac{m+r-2}{2}}$	$2^{m-r-1} + 2^{\frac{m-r-2}{2}}$
$2^{m-1} + 2^{\frac{m+r-2}{2}}$	$2^{m-r-1} - 2^{\frac{m-r-2}{2}}$

Table 1. Hamming weight and multiplicity in $\mathcal{C}$ when $p = 2$ and $m + r$ is even.

For $m = 5$, a 3-plateaued Boolean function and the corresponding binary linear code are given.

Example 3. Let $\Psi(x) = \zeta^{18}x^5 + \zeta^2 x^3$ be a mapping from $\mathbb{F}_{2^5}$ to $\mathbb{F}_{2^5}$, where $\mathbb{F}_{2^5}^\star = \langle\zeta\rangle$ with $\zeta^5 + \zeta^2 + 1 = 0$. Then, $\psi_1(x) = \mathrm{Tr}_2^{2^5}(\Psi(x))$ is the 3-plateaued Boolean function, and so the set $\mathcal{C}$ in (5) is a binary three-wight linear code with parameters $[31, 6]$, weight enumerator $1 + 59y^{16} + 3y^8 + 1y^{24}$ and weight distribution $(1, 59, 3, 1)$.

4.2 A new class of three-weight linear p-ary codes from weakly regular plateaued functions

In this subsection, we construct a new class of linear p-ary codes with few weights from weakly regular plateaued p-ary functions and determine their weight distributions. From now on, we assume that p is an odd prime and the function $\psi_1(x) = \mathrm{Tr}_p^{p^m}(\Psi(x))$ is weakly regular p-ary r-plateaued, where r is an integer with $0 \leq r \leq m$ and $\Psi : \mathbb{F}_{p^m} \to \mathbb{F}_{p^m}$ with $\Psi(0) = 0$. Let $\mathcal{C}$ be a linear p-ary code defined by (5) whose codewords are

denoted by $\tilde{c}_{\alpha,\beta}$. We first compute for all $\alpha \in \mathbb{F}_p$ and $\beta \in \mathbb{F}_{p^m}$, the Hamming weights of $\tilde{c}_{\alpha,\beta}$ and next determine the weight distribution of $\mathcal{C}$. By Proposition 2, if $\alpha = 0$, then we have $wt(\tilde{c}_{0,0}) = 0$ and $wt(\tilde{c}_{0,\beta}) = p^m - p^{m-1}$ for $\beta \neq 0$. For $\alpha \in \mathbb{F}_p^\star$, to compute $wt(\tilde{c}_{\alpha,\beta})$, we need the following.

Lemma 5. *Let $f : \mathbb{F}_{p^m} \to \mathbb{F}_p$ be r-plateaued, where r is an integer with $0 \leq r \leq m$. Define the sets $W := \{(\alpha,\beta) \in \mathbb{F}_p^\star \times \mathbb{F}_{p^m} \mid \widehat{\chi}_f(\alpha^{-1}\beta) = 0\}$ and*

$$WS := \{(\alpha,\beta) \in \mathbb{F}_p^\star \times \mathbb{F}_{p^m} \mid \widehat{\chi}_f(\alpha^{-1}\beta) \neq 0\}.$$

Then, the cardinalities of W and WS are equal respectively to $(p-1)(p^m - p^{m-r})$ and $(p-1)p^{m-r}$.

Proof. By Lemma 1, we have $\#\{\beta \in \mathbb{F}_{p^m} \mid \widehat{\chi_f}(\beta) = 0\} = p^m - p^{m-r}$ and $\#Supp(\widehat{\chi_f}) = p^{m-r}$, where $Supp(\widehat{\chi_f}) = \{\beta \in \mathbb{F}_{p^m} \mid \widehat{\chi_f}(\beta) \neq 0\}$. Hence, the result follows. □

For all $\alpha \in \mathbb{F}_p^\star$ and $\beta \in \mathbb{F}_{p^m}$, by Proposition 2, we have

$$wt(\tilde{c}_{\alpha,\beta}) = p^m - p^{m-1} - \frac{1}{p}\sum_{\omega \in \mathbb{F}_p^\star} \sigma_\omega\left(\sigma_\alpha(\widehat{\chi_{\psi_1}}(\alpha^{-1}\beta))\right). \tag{6}$$

Then there are two cases: $\widehat{\chi_{\psi_1}}(\alpha^{-1}\beta) = 0$ or $\neq 0$. If $(\alpha,\beta) \in W$, i.e., $\widehat{\chi_{\psi_1}}(\alpha^{-1}\beta) = 0$, then we have $wt(\tilde{c}_{\alpha,\beta}) = p^m - p^{m-1}$, that is, the number of codewords of Hamming weight $p^m - p^{m-1}$ is equal to the cardinality of W by Lemma 5. If $(\alpha,\beta) \in WS$, i.e., $\widehat{\chi_{\psi_1}}(\alpha^{-1}\beta) \neq 0$, to compute $wt(\tilde{c}_{\alpha,\beta})$ in (6), we use the following (see Lemma 3)

$$\widehat{\chi_{\psi_1}}(\alpha^{-1}\beta) = \epsilon\sqrt{p^*}^{m+r}\xi_p^{g(\alpha^{-1}\beta)},$$

where $\epsilon = \pm 1$, p^* denotes $\left(\frac{-1}{p}\right)p$ and g is a p-ary function over $Supp(\widehat{\chi_{\psi_1}})$. Notice that we have $\sigma_\alpha(\sqrt{p^*}^{m+r}) = \sigma_\alpha(\sqrt{p^*})^{m+r} = \left(\frac{\alpha}{p}\right)^{m+r}\sqrt{p^*}^{m+r}$, where σ_α is the automorphism of $\mathbb{Q}(\xi_p)$ for $\alpha \in \mathbb{F}_p^\star$. Then we get

$$\sigma_\omega\left(\sigma_\alpha(\widehat{\chi_{\psi_1}}(\alpha^{-1}\beta))\right) = \sigma_\omega\left(\epsilon\left(\frac{\alpha}{p}\right)^{m+r}\sqrt{p^*}^{m+r}\xi_p^{\alpha g(\alpha^{-1}\beta)}\right) =$$

$$\epsilon\left(\frac{\alpha}{p}\right)^{m+r}\sigma_\omega(\sqrt{p^*}^{m+r})\xi_p^{\omega\alpha g(\alpha^{-1}\beta)} = \epsilon\left(\frac{\alpha}{p}\right)^{m+r}\left(\frac{\omega}{p}\right)^{m+r}\sqrt{p^*}^{m+r}\xi_p^{\omega\alpha g(\alpha^{-1}\beta)},$$

where σ_ω is the automorphism of $\mathbb{Q}(\xi_p)$ for $\omega \in \mathbb{F}_p^\star$. Notice that $\left(\frac{a}{p}\right)^{m+r} = 1$ and $\sqrt{p^*}^{m+r} = \sqrt{p}^{m+r}$ if $m+r$ is even; otherwise, $\left(\frac{a}{p}\right)^{m+r} = \left(\frac{a}{p}\right)$ for $a \in \mathbb{F}_p^\star$. Hence, by (6) we have

$$wt(\tilde{c}_{\alpha,\beta}) = \begin{cases} p^m - p^{m-1} - \epsilon\frac{1}{p}\left(\frac{\alpha}{p}\right)\sqrt{p^*}^{m+r}\sum_{\omega\in\mathbb{F}_p^\star}\left(\frac{\omega}{p}\right)\xi_p^{\omega\alpha g(\alpha^{-1}\beta)}, & \text{if } m+r \text{ odd,} \\ p^m - p^{m-1} - \epsilon p^{\frac{m+r}{2}-1}\sum_{\omega\in\mathbb{F}_p^\star}\xi_p^{\omega\alpha g(\alpha^{-1}\beta)}, & \text{if } m+r \text{ even.} \end{cases}$$

We now investigate two cases. First, assume $m+r$ odd. If $g(\alpha^{-1}\beta) = 0$, then

$$wt(\tilde{c}_{\alpha,\beta}) = p^m - p^{m-1} - \epsilon\frac{1}{p}\left(\frac{\alpha}{p}\right)\sqrt{p^*}^{m+r}\sum_{\omega\in\mathbb{F}_p^\star}\left(\frac{\omega}{p}\right) = p^m - p^{m-1},$$

where we used $\sum_{\omega\in\mathbb{F}_p^\star}\left(\frac{\omega}{p}\right)=0$. If $g(\alpha^{-1}\beta)\neq 0$, then we have

$$\sum_{\omega\in\mathbb{F}_p^\star}\left(\frac{\omega}{p}\right)(\xi_p^{\omega})^{\alpha g(\alpha^{-1}\beta)}=\sigma_{\alpha g(\alpha^{-1}\beta)}\left(\sum_{\omega\in\mathbb{F}_p^\star}\left(\frac{\omega}{p}\right)\xi_p^{\omega}\right)=\sigma_{\alpha g(\alpha^{-1}\beta)}(\sqrt{p^*})=\left(\frac{\alpha g(\alpha^{-1}\beta)}{p}\right)\sqrt{p^*},$$

where we used $\sum_{\omega\in\mathbb{F}_p^\star}(\frac{\omega}{p})\xi_p^{\omega}=\sqrt{p^*}$. Hence,

$$\begin{aligned}wt(\tilde{c}_{\alpha,\beta})&=p^m-p^{m-1}-\epsilon\tfrac{1}{p}\sqrt{p^*}^{m+r+1}\left(\tfrac{\alpha^2}{p}\right)\left(\tfrac{g(\alpha^{-1}\beta)}{p}\right)\\&=p^m-p^{m-1}-\epsilon\tfrac{1}{p}\left(\tfrac{-1}{p}\right)^{\frac{m+r+1}{2}}p^{\frac{m+r+1}{2}}\left(\tfrac{g(\alpha^{-1}\beta)}{p}\right)\\&=p^m-p^{m-1}-\epsilon\,(-1)^{\frac{(p-1)(m+r+1)}{4}}p^{\frac{m+r-1}{2}}\left(\tfrac{g(\alpha^{-1}\beta)}{p}\right),\end{aligned}$$

where we used $\left(\frac{\alpha}{p}\right)\left(\frac{\alpha}{p}\right)=\left(\frac{\alpha^2}{p}\right)=1$, $p^*=\left(\frac{-1}{p}\right)p$ and $\left(\frac{-1}{p}\right)=(-1)^{\frac{p-1}{2}}$.
Now, assume $m+r$ even. If $g(\alpha^{-1}\beta)=0$, then we have

$$wt(\tilde{c}_{\alpha,\beta})=p^m-p^{m-1}-\epsilon p^{\frac{m+r-2}{2}}(p-1);$$

otherwise, we have $wt(\tilde{c}_{\alpha,\beta})=p^m-p^{m-1}+\epsilon p^{\frac{m+r-2}{2}}$ because if $g(\alpha^{-1}\beta)\neq 0$, then $\sum_{\omega\in\mathbb{F}_p^\star}\xi_p^{\alpha\omega g(\alpha^{-1}\beta)}=-1$ since $\sum_{j=0}^{p-1}x^j$ is the minimal polynomial of ξ_p over $\mathbb{Q}$.

We now collect in the following theorem the Hamming weights of the codewords of $\mathcal{C}$ defined by (5).

Theorem 2. *Let $\mathcal{C}$ be a linear p-ary code defined by (5). Assume that ψ_1 is weakly regular p-ary r-plateaued. Then, for all $\alpha\in\mathbb{F}_p$ and $\beta\in\mathbb{F}_{p^m}$, the Hamming weights of $\tilde{c}_{\alpha,\beta}$ are given as follows. For $\alpha=0$, we have $wt(\tilde{c}_{0,0})=0$ and $wt(\tilde{c}_{0,\beta})=p^m-p^{m-1}$ for $\beta\neq 0$. For $\alpha\in\mathbb{F}_p^\star$ and $\beta\in\mathbb{F}_{p^m}$, if $(\alpha,\beta)\in W$, i.e., $\widehat{\chi_{\psi_1}}(\alpha^{-1}\beta)=0$, then we get $wt(\tilde{c}_{\alpha,\beta})=p^m-p^{m-1}$, and if $(\alpha,\beta)\in WS$, i.e., $\widehat{\chi_{\psi_1}}(\alpha^{-1}\beta)\neq 0$, then*

- *when $m+r$ is odd,*

$$wt(\tilde{c}_{\alpha,\beta})=\begin{cases}p^m-p^{m-1}, & \text{if } \alpha\in\mathbb{F}_p^\star \text{ and } g(\alpha^{-1}\beta)=0,\\ p^m-p^{m-1}-\epsilon\,(-1)^{\frac{(p-1)(m+r+1)}{4}}p^{\frac{m+r-1}{2}}\left(\frac{g(\alpha^{-1}\beta)}{p}\right), & \text{if } \alpha, g(\alpha^{-1}\beta)\in\mathbb{F}_p^\star,\end{cases}$$

- *when $m+r$ is even,*

$$wt(\tilde{c}_{\alpha,\beta})=\begin{cases}p^m-p^{m-1}-\epsilon(p-1)p^{\frac{m+r-2}{2}}, & \text{if } \alpha\in\mathbb{F}_p^\star \text{ and } g(\alpha^{-1}\beta)=0,\\ p^m-p^{m-1}+\epsilon p^{\frac{m+r-2}{2}}, & \text{if } \alpha, g(\alpha^{-1}\beta)\in\mathbb{F}_p^\star,\end{cases}$$

where $\epsilon=\pm 1$ is the sign of $\widehat{\chi_{\psi_1}}$.

Now we are going to determine the weight distributions of the constructed codes given in Theorem 2. To do this, we first give the following result. By Lemma 4, the Walsh transform of g is written as

$$\widehat{\chi_g}(x)=\epsilon v p^{\frac{m-r}{2}}\xi_p^{\psi_1(-x)},$$

where $\epsilon = \pm 1$ denotes the sign of $\widehat{\chi_g}$ and $v \in \{1, i\}$ in $\mathbb{C}$. By using this for $x = 0$, we can compute the number of $b \in Supp(\widehat{\chi_{\psi_1}})$ such that $g(b) = j$ for all $j \in \mathbb{F}_p$. Set

$$N_g(j) := \#\{b \in Supp(\widehat{\chi_{\psi_1}}) \mid g(b) = j\}.$$

Notice that $g(b) = 0$ for all $b \notin Supp(\widehat{\chi_{\psi_1}})$, and $\#Supp(\widehat{\chi_{\psi_1}}) = p^{m-r}$. Hence, we have

$$\sum_{j=0}^{p-1} N_g(j) = p^{m-r}. \tag{7}$$

Remark 2. If g is balanced over $Supp(\widehat{\chi_{\psi_1}})$, we have $N_g(j) = p^{m-r-1}$ for all $j \in \mathbb{F}_p$.

We include the proof of the following proposition for making the paper self-contained (see, e.g., [5,10]).

Proposition 3. *We keep the above notations and assume that g is unbalanced over $Supp(\widehat{\chi_{\psi_1}})$. Then we have the following. If $m - r$ is even, then*

$$N_g(j) = \begin{cases} p^{m-r-1} + \epsilon p^{\frac{m-r-2}{2}}(p-1), j = 0, \\ p^{m-r-1} - \epsilon p^{\frac{m-r-2}{2}}, \qquad j \in \mathbb{F}_p^\star. \end{cases}$$

If $m - r$ is odd, then

$$N_g(j) = \begin{cases} p^{m-r-1}, & j = 0, \\ p^{m-r-1} + \epsilon p^{\frac{m-r-1}{2}} \left(\frac{j}{p}\right), j \in \mathbb{F}_p^\star, \end{cases}$$

where $\epsilon = \pm 1$ is the sign of $\widehat{\chi_g}$.

Proof. Using the Walsh value of unbalanced g at point zero, then we have

$$\widehat{\chi_g}(0) = \sum_{b \in Supp(\widehat{\chi_{\psi_1}})} \xi_p^{g(b)} = \sum_{j=0}^{p-1} N_g(j) \xi_p^j = \epsilon v p^{\frac{m-r}{2}} \xi_p^{\psi_1(0)}$$

equivalently,

$$\sum_{j=0}^{p-1} N_g(j) \xi_p^j - \epsilon v p^{\frac{m-r}{2}} = 0. \tag{8}$$

If $m - r$ is even, then $v = 1$. Because $\sum_{j=0}^{p-1} x^j$ is the minimal polynomial of ξ_p over the rational number field, then for all $j \in \mathbb{F}_p^\star$ we have

$$N_g(j) = a, \text{ and } N_g(0) = a + \epsilon p^{\frac{m-r}{2}}$$

for some constant a. By (7), $a + \epsilon p^{\frac{m-r}{2}} + (p-1)a = p^{m-r}$ from which one deduces $a = p^{m-r-1} - \epsilon p^{\frac{m-r}{2}-1}$.

If $m - r$ is odd, then $v = \begin{cases} 1, & \text{if } p \equiv 1 \pmod 4, \\ i, & \text{if } p \equiv 3 \pmod 4. \end{cases}$

Recall the well-known identity (see, e.g.,[9])

$$\sum_{j=0}^{p-1} \left(\frac{j}{p}\right) \xi_p^j = \begin{cases} p^{\frac{1}{2}}; & \text{if } p \equiv 1 \pmod 4, \\ i p^{\frac{1}{2}}, & \text{if } p \equiv 3 \pmod 4, \end{cases}$$

that is, $\sum_{j=0}^{p-1}\left(\frac{j}{p}\right)\xi_p^j = vp^{\frac{1}{2}}$. Thus, (8) can be rewritten as

$$\sum_{j=0}^{p-1} N_g(j)\xi_p^j - \epsilon p^{\frac{m-r-1}{2}} \sum_{j=0}^{p-1}\left(\frac{j}{p}\right)\xi_p^j = 0;$$

equivalently,

$$\sum_{j=0}^{p-1} \xi_p^j \left(N_g(j) - \epsilon p^{\frac{m-r-1}{2}}\left(\frac{j}{p}\right)\right) = 0.$$

Then for all $j \in \mathbb{F}_p^\star$, we have $N_g(j) = N_g(0) + \epsilon p^{\frac{m-r-1}{2}}\left(\frac{j}{p}\right)$. By (7), we obtain $\sum_{j=0}^{p-1} N_g(j) = pN_g(0) + \epsilon p^{\frac{m-r-1}{2}} \sum_{j=0}^{p-1}\left(\frac{j}{p}\right) = p^{m-r}$. Thus, since $\sum_{j=0}^{p-1}\left(\frac{j}{p}\right) = 0$, the proof is complete. □

We can derive from Remark 2 and Proposition 3 the weight distributions of the constructed codes.

Theorem 3. *Let $\mathcal{C}$ be a linear p-ary code defined by (5). Assume that ψ_1 is weakly regular p-ary r-plateaued and $m+r$ is even with $0 \le r \le m-2$ for $2 \le m$. Then, the Hamming weights of codewords and the weight distributions of $[p^m-1, m+1]$ code $\mathcal{C}$ are as in Tables 2 and 3 if g is unbalanced and balanced over $Supp(\widehat{\chi_{\psi_1}})$, respectively, where $\epsilon = \pm 1$ is the sign of $\widehat{\chi_{\psi_1}}$.*

Hamming weight w	Multiplicity A_w
0	1
$p^m - p^{m-1}$	$p^{m+1} - p^{m-r}(p-1) - 1$
$p^m - p^{m-1} - \epsilon(p-1)p^{\frac{m+r-2}{2}}$	$p^{m-r-1}(p-1) + \epsilon p^{\frac{m-r-2}{2}}(p-1)^2$
$p^m - p^{m-1} + \epsilon p^{\frac{m+r-2}{2}}$	$(p^{m-r} - p^{m-r-1})(p-1) - \epsilon p^{\frac{m-r-2}{2}}(p-1)^2$

Table 2. Hamming weight and multiplicity in $\mathcal{C}$ when $m+r$ is even and p is odd for unbalanced g

Hamming weight w	Multiplicity A_w
0	1
$p^m - p^{m-1}$	$p^{m+1} - p^{m-r}(p-1) - 1$
$p^m - p^{m-1} - \epsilon(p-1)p^{\frac{m+r-2}{2}}$	$p^{m-r-1}(p-1)$
$p^m - p^{m-1} + \epsilon p^{\frac{m+r-2}{2}}$	$\left(p^{m-r} - p^{m-r-1}\right)(p-1)$

Table 3. Hamming weight and multiplicity in $\mathcal{C}$ when $m+r$ is even and p is odd for balanced g

Proof. By Theorem 2, the numbers of codewords of Hamming weight 0 and of Hamming weight $p^m - p^{m-1}$ are equal respectively to 1 and $p^m - 1 + \#W = p^{m+1} + p^{m-r} - p^{m-r+1} - 1$. Now we are going to determine the weight distribution of $\mathcal{C}$ for $(\alpha, \beta) \in WS$, i.e., $\widehat{\chi_{\psi_1}}(\alpha^{-1}\beta) \neq 0$. Set

$$\begin{aligned} N_g(0) &:= \#\{\gamma \in Supp(\widehat{\chi_{\psi_1}}) \mid g(\gamma) = 0\}, \\ K_g(0) &:= \#\{(\alpha,\beta) \in \mathbb{F}_p^\star \times \mathbb{F}_{p^m} \mid g(\alpha^{-1}\beta) = 0\}, \\ KS_g &:= \#\{(\alpha,\beta) \in \mathbb{F}_p^\star \times \mathbb{F}_{p^m} \mid g(\alpha^{-1}\beta) \neq 0\}. \end{aligned}$$

Notice that for all $b \notin Supp(\widehat{\chi_{\psi_1}})$, $g(b) = 0$ by definition of g and so, $g(\alpha^{-1}\beta) = 0$ for all $(\alpha, \beta) \in W$. Hence, by Lemma 5, $K_g(0) = \#W + (p-1)N_g(0)$ and $KS_g = (p-1)p^m - K_g(0)$. Assume that g is unbalanced over $Supp(\widehat{\chi_{\psi_1}})$. Then, since $N_g(0) = p^{m-r-1} + \epsilon p^{(m-r-2)/2}(p-1)$ by Proposition 3, we have

$$K_g(0) = \#W + p^{m-r-1}(p-1) + \epsilon p^{\frac{m-r-2}{2}}(p-1)^2,$$

and $KS_g = (p^{m-r} - p^{m-r-1})(p-1) - \epsilon p^{(m-r-2)/2}(p-1)^2$. Hence, by Theorem 2, the numbers of codewords of Hamming weight $p^m - p^{m-1} - \epsilon(p-1)p^{(m+r-2)/2}$ and of Hamming weight $p^m - p^{m-1} + \epsilon p^{(m+r-2)/2}$ are equal to $K_g(0) - \#W$ and KS_g, respectively. Assume that g is balanced over $Supp(\widehat{\chi_{\psi_1}})$. By Remark 2, $N_g(0) = p^{m-r-1}$, and so we have $K_g(0) = \#W + p^{m-r-1}(p-1)$ and $KS_g = (p^{m-r} - p^{m-r-1})(p-1)$. As in the first case, the assertion holds. □

For $p = 3$ and $m = 3$, a weakly regular 3-ary 1-plateaued function and the corresponding linear 3-ary code are given as follows.

Example 4. Let $\Psi : \mathbb{F}_{3^3} \to \mathbb{F}_{3^3}$ be a map defined by $\Psi(x) = \zeta^{22}x^{13} + \zeta^7 x^4 + \zeta x^2$ where $\mathbb{F}_{3^3}^\star = \langle\zeta\rangle$ with $\zeta^3 + 2\zeta + 1 = 0$. A function $\psi_1(x) = \mathrm{Tr}_3^{3^3}(\Psi(x))$ is weakly regular 3-ary 1-plateaued with $\widehat{\chi_{\psi_1}}(b) \in \{0, -9\xi_3^{g(b)}\}$, where g is an unbalanced 3-ary function. Then, the set $\mathcal{C}$ in (5) is a three-wight linear 3-ary code with parameters $[26, 4]_3$, weight enumerator $1 + 62y^{18} + 2y^{24} + 16y^{15}$ and weight distribution $(1, 62, 2, 16)$.

Theorem 4. *Let $\mathcal{C}$ be a linear p-ary code defined by (5). Assume that ψ_1 is weakly regular p-ary r-plateaued and $m + r$ is odd with $0 \le r \le m - 1$. Then, the Hamming weights of codewords and the weight distributions of $[p^m - 1, m + 1]$ code $\mathcal{C}$ are as in Tables 4 and 5 if g is unbalanced and balanced over $Supp(\widehat{\chi_{\psi_1}})$, respectively, where $\epsilon = \pm 1$ is the sign of $\widehat{\chi_{\psi_1}}$.*

Hamming weight w	Multiplicity A_w
0	1
$p^m - p^{m-1}$	$p^{m+1} - p^{m-r-1}(p-1)^2 - 1$
$p^m - p^{m-1} - \epsilon\,(-1)^{\frac{(p-1)(m+r+1)}{4}}\,p^{\frac{m+r-1}{2}}$	$\frac{1}{2}(p^{m-r-1} + \epsilon p^{\frac{m-r-1}{2}})(p-1)^2$
$p^m - p^{m-1} + \epsilon\,(-1)^{\frac{(p-1)(m+r+1)}{4}}\,p^{\frac{m+r-1}{2}}$	$\frac{1}{2}(p^{m-r-1} - \epsilon p^{\frac{m-r-1}{2}})(p-1)^2$

Table 4. Hamming weight and multiplicity in $\mathcal{C}$ when $m + r$ and p are odd for unbalanced g

Hamming weight w	Multiplicity A_w
0	1
$p^m - p^{m-1}$	$p^{m+1} - p^{m-r-1}(p-1)^2 - 1$
$p^m - p^{m-1} - \epsilon\,(-1)^{\frac{(p-1)(m+r+1)}{4}}\,p^{\frac{m+r-1}{2}}$	$\frac{1}{2}p^{m-r-1}(p-1)^2$
$p^m - p^{m-1} + \epsilon\,(-1)^{\frac{(p-1)(m+r+1)}{4}}\,p^{\frac{m+r-1}{2}}$	$\frac{1}{2}p^{m-r-1}(p-1)^2$

Table 5. Hamming weight and multiplicity in $\mathcal{C}$ when $m + r$ and p are odd for balanced g

Proof. Set $N_g(j) := \#\{\gamma \in Supp(\widehat{\chi_{\psi_1}}) \mid g(\gamma) = j\}$ and $K_g(j) := \#\{(\alpha, \beta) \in \mathbb{F}_p^\star \times \mathbb{F}_{p^m} \mid g(\alpha^{-1}\beta) = j\}$ for all $j \in \mathbb{F}_p$. Notice that for all $b \notin Supp(\widehat{\chi_{\psi_1}})$, $g(b) = 0$ by

definition of g and so, $g(\alpha^{-1}\beta) = 0$ for all $(\alpha, \beta) \in W$. Then, by Lemma 5, $K_g(0) = \#W + (p-1)N_g(0)$ where $N_g(0) = p^{m-r-1}$ (see Remark 2 and Proposition 3). Hence, by Theorem 2, the number of codewords of Hamming weight $p^m - p^{m-1}$ is equal to $p^m - 1 + K_g(0) = p^{m+1} + 2p^{m-r} - p^{m-r+1} - p^{m-r-1} - 1$. Moreover, the number of codewords of Hamming weight $p^m - p^{m-1} - \epsilon\,(-1)^{(p-1)(m+r+1)/4}\,p^{(m+r-1)/2}$ and of Hamming weight $p^m - p^{m-1} + \epsilon\,(-1)^{(p-1)(m+r+1)/4}\,p^{(m+r-1)/2}$ is equal respectively to $\sum_{j\in\{1,\ldots,p-1\},\left(\frac{j}{p}\right)=1}(p-1)N_g(j)$ and $\sum_{j\in\{1,\ldots,p-1\},\left(\frac{j}{p}\right)=-1}(p-1)N_g(j)$. If g is unbalanced, then by Proposition 3,

$$\begin{aligned}\sum_{j\in\{1,\ldots,p-1\},\left(\frac{j}{p}\right)=1}(p-1)N_g(j) &= \sum_{j\in\{1,\ldots,p-1\},\left(\frac{j}{p}\right)=1}(p-1)(p^{m-r-1} + \epsilon p^{\frac{m-r-1}{2}}) \\ &= \tfrac{(p-1)^2}{2}(p^{m-r-1} + \epsilon p^{\frac{m-r-1}{2}})\end{aligned}$$

and

$$\begin{aligned}\sum_{j\in\{1,\ldots,p-1\},\left(\frac{j}{p}\right)=-1}(p-1)N_g(j) &= \sum_{j\in\{1,\ldots,p-1\},\left(\frac{j}{p}\right)=-1}(p-1)(p^{m-r-1} - \epsilon p^{\frac{m-r-1}{2}}) \\ &= \tfrac{(p-1)^2}{2}(p^{m-r-1} - \epsilon p^{\frac{m-r-1}{2}}).\end{aligned}$$

If g is balanced then by Remark 2, $\sum_{j\in\{1,\ldots,p-1\},\left(\frac{j}{p}\right)=1}(p-1)N_g(j) = \dfrac{(p-1)^2}{2}p^{m-r-1}$ and $\sum_{j\in\{1,\ldots,p-1\},\left(\frac{j}{p}\right)=-1}(p-1)N_g(j) = \dfrac{(p-1)^2}{2}p^{m-r-1}$. The proof is complete. □

Remark 3. We finally should remark that if we assume only the weakly regular bentness in this paper, then we can recover the results given in [10] by Mesnager. Therefore, this paper can be viewed as an extension of [10] to the notion of weakly regular r-plateaued functions for any positive integer r.

5 The constructed three-weight linear codes for secret sharing schemes

In this section, we consider our linear codes presented in Section 4 for secret sharing schemes. A linear code provides a pair of secret sharing schemes, based on a linear code $\mathcal{C}$ and its dual code $\mathcal{C}^\perp$. For the secret sharing scheme based on the dual code $\mathcal{C}^\perp$, we need to find all minimal codewords of $\mathcal{C}$. We say that a vector $\tilde{a}$ covers a vector $\tilde{b}$ if $supp(\tilde{b}) \subset supp(\tilde{a})$. Then, if a nonzero codeword $\tilde{a}$ of $\mathcal{C}$ does not cover any other nonzero codeword of $\mathcal{C}$, then $\tilde{a}$ is called *minimal codeword* of $\mathcal{C}$. The *covering problem* is to find all the minimal codewords of $\mathcal{C}$. In general, this problem is very hard, but it can be easy for some linear codes. Then the main question is how to find a linear code whose all nonzero codewords are minimal. For more details, we send the reader to [4].

Lemma 6. *[1] Let $\mathcal{C}$ be a linear code over $\mathbb{F}_p$. Every nonzero codeword of $\mathcal{C}$ is minimal if $\frac{p-1}{p} < \frac{w_{min}}{w_{max}}$, where w_{min} and w_{max} denote the minimum and maximum nonzero weights in $\mathcal{C}$, respectively.*

We now consider the constructed linear codes in Theorems 1, 3 and 4. Let $\mathcal{C}$ be the binary linear code of Theorem 1 and $m+r$ be even. Then we readily see that $\frac{1}{2} < \frac{w_{min}}{w_{max}}$, where $w_{min} = 2^{m-1} - 2^{(m+r-2)/2}$ and $w_{max} = 2^{m-1} + 2^{(m+r-2)/2}$ since we have $3 \cdot 2^{(m+r)/2} < 2^m$ for $m \geq 4$ and $0 \leq r \leq m-4$. Hence, by Lemma 6, all nonzero codewords of $\mathcal{C}$ given in Theorem 1 are minimal if $m \geq 4$ and $0 \leq r \leq m-4$.

Let p be any odd prime, $m+r$ be even and $\mathcal{C}$ be the linear p-ary code of Theorem 3. If $\epsilon = 1$, we have $w_{min} = p^m - p^{m-1} - (p-1)p^{(m+r-2)/2}$ and $w_{max} = p^m - p^{m-1} + p^{(m+r-2)/2}$. If $\epsilon = -1$, then $w_{min} = p^m - p^{m-1} - p^{(m+r-2)/2}$ and $w_{max} = p^m - p^{m-1} + (p-1)p^{(m+r-2)/2}$. For both cases, we see that $\frac{p-1}{p} < \frac{w_{min}}{w_{max}}$ for $m \geq 4$ and $0 \leq r \leq m-4$ since we have $(p+1)p^{(m+r)/2} < p^m$ if $\epsilon = 1$ and $(p^2-p+1)p^{(m+r)/2} < p^m(p-1)$ if $\epsilon = -1$. Hence, by Lemma 6, all nonzero codewords of $\mathcal{C}$ given in Theorem 3 are minimal if $m \geq 4$ and $0 \leq r \leq m-4$.

Let p be any odd prime, $m+r$ be odd and $\mathcal{C}$ be the linear p-ary code of Theorem 4. Then we see that $\frac{p-1}{p} < \frac{w_{min}}{w_{max}}$, where $w_{min} = p^m - p^{m-1} - p^{(m+r-1)/2}$ and $w_{max} = p^m - p^{m-1} + p^{(m+r-1)/2}$ since we have $(2p-1)p^{(m+r+1)/2} < p^m(p-1)$ for $m \geq 3$ and $0 \leq r \leq m-3$. Hence, by Lemma 6, all nonzero codewords of $\mathcal{C}$ given in Theorem 4 are minimal if $m \geq 3$ and $0 \leq r \leq m-3$.

6 Conclusion

The paper studies for the first time constructions of linear codes with few weights from weakly regular plateaued functions. We first present a new family of binary three-weight linear codes from plateaued Boolean functions and their weight distributions. In odd characteristic p, we introduce the notion of (weakly) regular plateaued functions and give concrete examples of these functions. We next present a new family of three-weight linear p-ary codes from weakly regular plateaued functions, and their weight distributions. We finally analyse the constructed linear codes in this paper for secret sharing schemes. The constructed linear codes are inequivalent to the known ones (since there is no code with the obtained parameters) in literature as far as we know.

Acknowledgment

The third author is supported by TÜBİTAK (the Scientific and Technological Research Council of Turkey), program no: BİDEB 2214/A.

References

1. A. Ashikhmin, Variations on minimal codewords in linear codes. In Proc. 11th Int. Symp. AAECC, Springer Berlin Heidelberg, 96-105 (1995)
2. C. Ding.: A construction of binary linear codes from Boolean functions. Discrete mathematics. Vol. 339, No. 9, pp. 2288-2303 (2016)
3. K. Ding., C. Ding.: Binary linear codes with three weights. IEEE Communications Letters. Vol. 18, No. 11, pp. 1879-1882 (2014)
4. K. Ding., C. Ding.: A class of two-weight and three-weight codes and their applications in secret sharing. IEEE T INFORM THEORY. Vol. 61, No. 11, pp. 5835-5842 (2015)

5. T. Helleseth., A. Kholosha.: Bent functions and their connections to combinatorics. Surveys in Combinatorics, Cambridge University Press, pp. 91-126, (2013)
6. T. Helleseth., A. Kholosha.: Monomial and quadratic bent functions over the finite fields of odd characteristic. IEEE T INFORM THEORY, Vol. 52, No. 5, pp. 2018- 2032 (2006)
7. W. C. Huffman, V. Pless. Fundamentals of error-correcting codes. Cambridge, U.K.: Cambridge University press, (2010)
8. J.Y. Hyun, J. Lee, Y. Lee: Explicit criteria for construction of plateaued functions. IEEE T INFORM THEORY, Vol. 62, No. 12, pp. 7555 - 7565 (2016)
9. R. Lidl, H. Niederreiter.: Finite fields. 2nd Edition, Cambridge University Press (1997)
10. S. Mesnager.: Linear codes with few weights from weakly regular bent functions based on a generic construction. Cryptography and Communications, Vol. 9, No. 1, pp. 71-84 (2017)
11. S. Mesnager, F. Özbudak, A. Sınak: Results on Characterizations of Plateaued Functions in Arbitrary Characteristic. Cryptography and Information Security in the Balkans, BalkanCryptSec 2015, Koper, Slovenia, Revised Selected Papers, Pasalic, E., Knudsen, L.R.(eds.) LNCS 9540, Springer. 17–30 (2016)
12. C. Tang., N. Li., Y. Qi., Z. Zhou., T. Helleseth.: Linear codes with two or three weights from weakly regular bent functions. IEEE T INFORM THEORY. Vol. 62, No. 3, pp. 1166-1176 (2016)
13. K. Ireland., M. Rosen.: A Classical introduction to modern number theory, 2nd ed. New York: Springer-Verlag, Vol. 84, Graduate Texts in Mathematics (1990)
14. G. Xu., X. Cao.: Linear codes with two or three weights from some functions with low Walsh spectrum in odd characteristic. arXiv:1510.01031 (2015)
15. Y. Zheng, X.M. Zhang: Plateaued functions. Information and Communication Security. Springer, Berlin Heidelberg. pp. 284-300 (1999)
16. Z. Zhou., N. Li., C. Fan., T. Helleseth.: Linear codes with two or three weights from quadratic bent functions. DESIGN CODE CRYPTOGR. Vol. 81, No. 2, pp. 283-295 (2016)